

Bernoulli potential at a superconductor surface

Pavel Lipavský, Jan Koláček and Jiří J. Mareš

Institute of Physics, Academy of Sciences, Cukrovarnická 10, 16258 Praha 6, Czech Republic

Klaus Morawetz

Max-Planck-Institute for the Physics of Complex Systems, Noethnitzer Str. 38, 01187 Dresden, Germany

The electrostatic Bernoulli potential measured at the surface of a superconductor via Kelvin capacitive coupling is shown to be independent of the pairing mechanism. This contrasts with the Bernoulli potential in the bulk where contributions due to pairing dominate close to T_c .

In 1968 Bok and Klein¹ have first observed non-zero Hall voltage in the Meissner state of a type-I superconductor. Instead of Ohmic contacts, they have used the Kelvin capacitive coupling between the surface and the electrodes. Similar measurements have been performed by Brown and Morris^{2,3} and more recently by Chiang and Shevchenko^{4,5}.

Among these experiments, the highest accuracy has been achieved by Morris and Brown³. They found a perfect agreement with the prediction of van Vijfeijken and Staas⁶ in which the original Bernoulli potential (derived by Bopp⁷ and made familiar by London⁸) is extended by the fountain term of the two-fluid model. In particular, this measurement shows no traces of effects predicted later within more sophisticated thermodynamical approaches^{9–11}.

In this paper we show that the agreement of the two-fluid theory with experiment is in some sense fortuitous. This theory has been developed for the electrostatic potential inside the superconductor but it is the surface potential that is measured by the Kelvin method. The surface potential is influenced by the surface dipole which brings a step-like contribution to the electrostatic potential^{12,13}. With the surface dipole included, the measured surface potential agrees with the prediction of the thermodynamic approaches.

The Hall voltage in superconductors is a difference between electrostatic potentials, φ , at the contacts. All theories agree that this potential has the form of the Bernoulli potential,

$$e\varphi \propto -\frac{1}{2}mv^2, \quad (1)$$

where v is the velocity of the condensate and m is the mass of electrons. The point of disagreement concerns the amplitude of this potential.

London⁸ assumed that the electric force balances the inertial and Lorentz forces. The energy-conserving integral of motion then yields the Bernoulli potential,

$$e\varphi = -\frac{1}{2}mv^2. \quad (2)$$

This formula applies only at zero temperature. To cover finite temperatures, van Vijfeijken and Staas⁶ took into account the fact that the normal electrons feel the electric

field. Since they stay at rest, there must be a balancing force from the interaction with the condensate. The reaction force acts on the condensate and adds to the electric force. This picture leads to the electrostatic potential which is proportional to the fraction of superconducting electrons,

$$e\varphi = -\frac{n_s}{n}\frac{1}{2}mv^2. \quad (3)$$

The formula (3) represents the level of theory before the first successful experiment appeared. With respect to experiments it is profitable to express the Bernoulli potential (3) as a function of the magnetic field B . At the surface, $v = \frac{e}{m}\lambda B$, where $\lambda = \sqrt{\frac{m}{e^2 n_s \mu}}$ is the London penetration depth, therefore the two-fluid potential (3) reads

$$e\varphi = -\frac{1}{n}\frac{B^2}{2\mu}. \quad (4)$$

Morris and Brown³ experimentally confirmed the validity of (4) in a wide temperature range including the vicinity of the critical temperature.

As discussed by Bok and Klein¹, there is also a simple theoretical argument supporting formula (4). In the absence of dissipation, the only force acting on the lattice is due to the electric field, $-\nabla\varphi$. The total force per unit area acting on a superconducting slab,

$$F = \int_{\text{right}}^{\text{left}} dx en \nabla\varphi = en\varphi_{\text{left}} - en\varphi_{\text{right}}, \quad (5)$$

is proportional to the difference of the surface potentials. Here, $-en$ stands for the charge density of the ionic lattice. In the magnetic field B , the slab with the current density J [in A/m] feels the Lorentz force $F = BJ$. Using the Ampere rule, $B_{\text{right}} - B_{\text{left}} = \mu J$, and the mean magnetic field, $B = \frac{1}{2}(B_{\text{right}} + B_{\text{left}})$, the Lorentz force can be expressed as the difference between the left and right magnetic pressures,

$$F = BJ = \frac{B_{\text{right}}^2}{2\mu} - \frac{B_{\text{left}}^2}{2\mu}. \quad (6)$$

One can see that the Lorentz force (6) agrees with (5) for potential (4).

In spite of its internal consistency and the satisfactory agreement with the experimental data of Bok and Klein, the two-fluid theory came under suspicion when studies^{9–11} based on thermodynamic relations and BCS analyses have been accomplished. These approaches result in a potential

$$e\varphi = -\frac{\partial n_s}{\partial n} \frac{1}{2} m v^2. \quad (7)$$

For the phenomenological density of condensate,

$$n_s = n \left(1 - \frac{T^4}{T_c^4} \right), \quad (8)$$

potential (7) reads

$$e\varphi = -\frac{n_s}{n} \frac{1}{2} m v^2 + 4 \frac{n_n}{n} \frac{\partial \ln T_c}{\partial \ln n} \frac{1}{2} m v^2, \quad (9)$$

where $n_n = n - n_s$ is the density of normal electrons. The first term is identical to the two-fluid potential (3), the second is the difference which we will call the thermodynamic correction.

Formulas (7) and (3), are very different close to T_c . Within the BCS relations and the parabolic band approximation, one can rearrange (9) as

$$e\varphi = -\frac{n_s}{n} \frac{1}{2} m v^2 - \frac{4}{3} \frac{n_n}{n} \ln \frac{T_D}{T_c} \frac{1}{2} m v^2. \quad (10)$$

Since the Debye temperature T_D is much larger than the critical temperature T_c , the second term dominates at higher temperatures, approximately for $T > \frac{2}{3} T_c$.

The thermodynamic correction arises from the pairing mechanism. According to the BCS critical temperature, $T_c \approx T_D e^{-1/\mathcal{D}V}$, where $\mathcal{D}$ is the single-spin density of states and V is the BCS interaction parameter, the potential (9) can be written as

$$e\varphi = -\frac{n_s}{n} \frac{1}{2} m v^2 - \frac{n_n}{n} \frac{1}{\mathcal{D}V} \frac{2}{3} m v^2. \quad (11)$$

As speculated by Rickayzen¹⁰, the Hall voltage at temperatures close to T_c can offer an experimental access to the BCS interaction V . Rickayzen expressed his surprise that Bok and Klein¹ did not observe any trace of the thermodynamic correction at the temperature $T = 4.2$ K for lead with $T_c = 7.2$ K. For $T_D = 105$ K the factors in (10) are comparable, $\frac{n_s}{n} = 0.88$ and $\frac{4}{3} \frac{n_n}{n} \ln \frac{T_D}{T_c} = 0.42$.

While the measurement of Bok and Klein¹ was not sufficiently precise and the thermodynamic correction could be lost in the experimental error, later measurement of Morris and Brown³ clearly proved that the Hall voltage is independent of temperature. They also used lead and explored the temperature range from 4.2 to 7.0 K. At $T = 7$ K the thermodynamic correction dominates the potential, since the factors in (10) are $\frac{n_s}{n} = 0.1$ and

$\frac{4}{3} \frac{n_n}{n} \ln \frac{T_D}{T_c} = 3.2$. If present, the thermodynamic correction would amplify the signal thirty times. This is far beyond the experimental error, estimated to be less than 10%. Briefly, the thermodynamic correction turned out to be absent. It should be mentioned that Morris and Brown expected a non-zero thermodynamic correction indicated by their less precise preliminary measurement² at $T = 4.2$ K.

One might feel that the experimental results imply that the thermodynamic correction is an ill-defined concept. It is, however, not the case. The electrostatic potential given by (7) applies in the bulk. To obtain the potential at the surface, one cannot simply extrapolate this bulk value. At the surface there is a surface dipole which leads to a discontinuity in the potential.^{12,13}

The surface dipole in a metal introduces a step of several eV's in the electrostatic potential. In the jellium model this step equals to the Fermi energy E_F . With the exchange and correlation energy included, the surface dipole is reduced by half which shows the importance of correlations. The phase transition into the superconducting state increases the correlation energy by the condensation energy. The surface dipole of the superconductor is thus slightly smaller than the dipole of the same crystal in the normal state. This difference is of the order of Δ^2/eE_F .

When a supercurrent flows close to the surface, the condensation energy is reduced. The surface dipole thus slightly increases, reaching the normal state value at the critical current. This current dependent part of the surface dipole contributes to the potential observed by the capacitive coupling^{1–3}. Now we show that the surface dipole cancels the thermodynamic contribution (second term of (9)) to the Hall voltage.

In our formulation we follow Rickayzen¹⁰. The increase of the free energy due to the current equals the kinetic energy of the condensate,

$$\delta\mathcal{F} = n_s \frac{1}{2} m v^2. \quad (12)$$

The electrochemical potential, $\nu = E_F + \delta\nu_v + e\varphi$, is constant in the whole system, therefore $e\varphi = -\delta\nu_v$. Since $\nu = \frac{\partial\mathcal{F}}{\partial n}$, the velocity variation of the local chemical potential is $\delta\nu_v = \frac{\partial}{\partial n} \delta\mathcal{F}$. Accordingly, the electrostatic potential induced by the current is given by

$$e\varphi = -\frac{\partial}{\partial n} \delta\mathcal{F}. \quad (13)$$

With $\delta\mathcal{F}$ given by (12), from (13) one recovers (7).

The step of the electrostatic potential at the surface is given by the Budd-Vannimenus theorem¹³,

$$e\varphi_{\text{surf}} - e\varphi = n \frac{\partial}{\partial n} \frac{\delta\mathcal{F}}{n}. \quad (14)$$

We note that Budd-Vannimenus theorem has been derived for the surface potential of the normal metal. Formula (14) is a modification restricted to the potential caused by the supercurrent.

From (14) and the bulk potential (13) one finds the electrostatic potential at the surface,

$$e\varphi_{\text{surf}} = -\frac{n_s}{n}\frac{1}{2}mv^2. \quad (15)$$

This potential is identical to the experimentally confirmed two-fluid potential (3). Accordingly the surface dipole cancels exactly the thermodynamic corrections.

In conclusion, the electrostatic potential observed at the surface of superconductors is determined by the magnetic pressure. The only material parameter accessible by its measurement is the charge density of the ionic lattice. In the bulk, the electrostatic potential depends also on the pairing mechanism.

Since the thermodynamic corrections cannot be measured on the surface, it is desirable to observe the internal electric field directly in the bulk of a superconductor. A new experiment in this direction has been reported by Kumagai *et al*¹⁴ who have measured the effect of charge density deviations on the nuclear quadrupole resonance.

It is not obvious how the surface potential works in layered structures of high- T_c materials. A measurement of the Hall voltage on these materials, however, involves enormous technical difficulties.¹⁵

The authors thank Nai Kwong for helpful discussions. This work was supported by GAČR 202000643 and 202990410, GAAV A1010806 and A1010919 grants. The European ESF program VORTEX is also acknowledged.

-
- ¹ J. Bok and J. Klein, Phys. Rev. Lett. **20**, 660 (1968).
 - ² J. B. Brown and T. D. Morris, Proc. 11th Int. Conf. Low. Temp. Phys., Vol. 2, 768 (St. Andrews, 1968).
 - ³ T. D. Morris and J. B. Brown, Physica **55**, 760 (1971).
 - ⁴ Yu. N. Chiang and O. G. Shevchenko, Fiz. Nizk. Temp. **12**, 816 (1986) [Sov. J. Low. Temp. Phys. **12**, 462 (1986)].
 - ⁵ Yu. N. Chiang and O. G. Shevchenko, Fiz. Nizk. Temp. **22**, 669 (1996) [Sov. J. Low. Temp. Phys. **22**, 513 (1996)].
 - ⁶ A. G. van Vijfeijken and F. S. Staas, Phys. Lett. **12**, 175 (1964).
 - ⁷ F. Bopp, Z. f. Phys. **107**, 623 (1937).
 - ⁸ F. London, *Superfluids* (Willey, New York, 1950), Vol. I, Sec. 8.
 - ⁹ C. J. Adkins and J. R. Waldram, Phys. Rev. Lett. **21**, 76 (1968).
 - ¹⁰ G. Rickayzen, J. Phys. C **2**, 1334 (1969).
 - ¹¹ K. M. Hong, Phys. Rev. B **12**, 1766 (1975).
 - ¹² J. R. Waldram, *Superconductivity of Metals and Cuprates*, Chap. 2.3, (Arrowsmith, Bristol 1996).
 - ¹³ A. Kleina and K. F. Wojciechowski, *Metal Surface Electron Physics*, (Elsevier Sciencies, Oxford, 1996).
 - ¹⁴ K. Kumagai, K. Nozaki and Y. Matsuda, Phys. Rev. B **63**, 144502 (2001).
 - ¹⁵ Tom Lemberger, private communication.